arXiv:1211.6685v1 [hep-th] 28 Nov 2012

Hawking Radiation in a Plebański-Demiański Black Hole

Jesús A. Cázares*

Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN
P.O.Box 14-740, 07000 México D.F., México and
Rudjer Bošković Institute, P.O.Box 180, HR-10002 Zagreb, Croatia

November 29, 2012

Abstract

We show in this paper the flux of Hawking radiation in a Plebański-Demiański black hole from gauge and gravitational anomalies point of view. We also show this calculations are general and the results from earlier known articles can be obtained from this one.

1 Introduction

The black hole radiation also called Hawking radiation, was originally reported by Zeldovich and Starobinsky in 1971 [1] and it is one of the more interesting known effect. This effect is a consequence of the combination of two modern theories: quantum mechanics and general relativity; because arise from quantum fileds in a background space–time with an event horizont, such like a black hole. It is founded that the occupation number spectrum of quantum filled modes in the vacuum state corresponds with the blackbody at fixed temperature given by the surface gravity of the horizon.

There are several explanation of Hawking radiation. The derivation by Hawking [2, 3] is direct and physical, he calculates the Bogoliubov coefficients between the in and out states of fields in the black hole background. The derivation based in quantum gravity [4] is fast and elegant, but need microscopic foundation. The derivation based in string theory [5, 6] has logical consistence foundation but can be applied only to special solutions and do not explain the simplicity and generality of the results inferred from other methods.

Robinson and Wilczek [7] have recently showed a procedure for Hawking radiation via gravitational anomalies cancelation in a Schwarzschild black hole metric. This original idea soon was extended to Reissner-Nördstrom metric [8] to rotating black holes [9, 10, 11, 12] and even to metrics with NUT parameter [13]. Opening at this form the important question: How is Hawking radiation for the most general black hole?

In the present article, we answer that question for the most general D–type metric (Plebański-Demiański metric) and show the form of fluxes and the Hawking temperature. The article is organized at this form. In the second section we introduce the Plebański-Demiański metric and we get the effective action, corresponding to $(1+1)$–dimensional metric in a dilaton background and a gauge field. In sections 3 and 4, we make the calculation of the gauge and gravitational anomalies that arise the Hwking radiation at the event horizon and cosmological horizon, respectively. We show two cases in section 5 like limit cases: the Kerr-Newman-de Sitter black hole metric and NUT-Kerr-Newman-de Sitter and we compare with the results obtained in the literature [11, 13]. Finally, the conclusions are showed in section 6.

---

[*] cazares@irb.hr

2 The Effective action of Plebański-Demiański black hole

As it is known, the most general D–type metric is the Plebański-Demiański metric [14]

$$
\begin{aligned}
\mathrm{d}s^2 &= \frac{1}{(1-\hat{p}\hat{r})^2}\left[\frac{\mathcal{Q}(\mathrm{d}\hat{\tau}-\hat{p}^2\mathrm{d}\hat{\sigma})^2}{\hat{r}^2+\hat{p}^2}-\frac{\mathcal{P}(\mathrm{d}\hat{\tau}+\hat{r}^2\mathrm{d}\hat{\sigma})^2}{\hat{r}^2+\hat{p}^2}-(\hat{r}^2+\hat{p}^2)\Big(\frac{\mathrm{d}\hat{p}^2}{\mathcal{P}}+\frac{\mathrm{d}\hat{r}^2}{\mathcal{Q}}\Big)\right], \qquad (1)\\
\mathcal{A} &= \frac{\hat{e}+\mathrm{i}\hat{g}}{\hat{r}+\mathrm{i}\hat{p}}\Big(\mathrm{d}\hat{\tau}-\mathrm{i}\hat{p}\hat{r}\mathrm{d}\hat{\sigma}\Big), \qquad (2)
\end{aligned}
$$

where

$$
\begin{aligned}
\mathcal{P} &= \hat{k}+2\hat{n}\hat{p}-\hat{\epsilon}\hat{p}^2+2\hat{m}\hat{p}^3-\left(\hat{k}+\hat{e}^2+\hat{g}^2+\frac{\Lambda}{3}\right)\hat{p}^4, \qquad (3)\\
\mathcal{Q} &= (\hat{k}+\hat{e}^2+\hat{g}^2)-2\hat{m}\hat{r}+\hat{\epsilon}\hat{r}^2-2\hat{n}\hat{r}^3-\left(\hat{k}+\frac{\Lambda}{3}\right)\hat{r}^4, \qquad (4)
\end{aligned}
$$

and $\hat{m}$, $\hat{n}$, $\hat{e}$, $\hat{g}$, $\hat{\epsilon}$, $\hat{k}$, Λ are arbitrary real parameters. Recently has been showed that it is possible to put this metric in the form [15]

$$
\begin{aligned}
\mathrm{d}s^2 &= \frac{1}{\Omega^2}\left\{\frac{\hat{\mathcal{Q}}}{\rho^2}\left[\mathrm{d}t-\frac{a\sin^2\theta+4l\sin^2\frac{\theta}{2}}{\Xi}\mathrm{d}\varphi\right]^2-\frac{\rho^2}{\hat{\mathcal{Q}}}\mathrm{d}r^2-\frac{\hat{\mathcal{P}}}{\rho^2}\left[a\mathrm{d}t-\frac{r^2+(a+l)^2}{\Xi}\mathrm{d}\varphi\right]^2-\frac{\rho^2}{\hat{\mathcal{P}}}\sin^2\theta\mathrm{d}\theta^2\right\}, \qquad (5)\\
\mathcal{A} &= A_\mu+\mathrm{i}B_\mu \qquad (6)\\
&= \left(\frac{Qr+G(l+a\cos\theta)}{r^2+(l+a\cos\theta)^2},0,0,-\frac{Qr\Big(4al\sin^2\frac{\theta}{2}+a^2\sin^2\theta\Big)+G(l+a\cos\theta)\Big(r^2+(l+a)^2\Big)}{a\Xi\Big(r^2+(l+a\cos\theta)^2\Big)}\right)\\
&\quad +\mathrm{i}\left(\frac{Gr-Q(l+a\cos\theta)}{r^2+(l+a\cos\theta)^2},0,0,\frac{-Gr\Big(4al\sin^2\frac{\theta}{2}+a^2\sin^2\theta\Big)+Q(l+a\cos\theta)\Big(r^2+(l+a)^2\Big)}{a\Xi\Big(r^2+(l+a\cos\theta)^2\Big)}\right); \qquad (7)
\end{aligned}
$$

where A_μ and B_μ are the electric and magnetic potentials and

$$
\begin{aligned}
\Omega &= 1-\frac{\alpha}{\omega}(l+a\cos\theta)r, \qquad (8)\\
\rho^2 &= r^2+(l+a\cos\theta)^2, \qquad (9)\\
\Xi &= 1+\frac{\Lambda}{3}(a^2-l^2), \qquad (10)\\
\hat{\mathcal{P}} &= \sin^2\theta(1-a_3\cos\theta-a_4\cos^2), \qquad (11)\\
\hat{\mathcal{Q}} &= (\omega^2k+Q^2+G^2)-2Mr+\epsilon r^2-2\alpha\frac{n}{\omega}r^3-\left(\alpha^2k+\frac{\Lambda}{3}\right)r^4, \qquad (12)\\
a_3 &= 2\alpha\frac{a}{\omega}m-4\alpha^2\frac{al}{\omega^2}(\omega^2k+Q^2+G^2)-4\frac{\Lambda}{3}al, \qquad (13)\\
a_4 &= -\alpha^2\frac{a^2}{\omega^2}(\omega^2k+Q^2+G^2)-\frac{\Lambda}{3}a^2, \qquad (14)\\
\epsilon &= \frac{\omega^2k}{a^2-l^2}+4\alpha\frac{l}{\omega}m-(a^2+3l^2)\left(\frac{\alpha^2}{\omega^2}(\omega^2k+Q^2+G^2)+\frac{\Lambda}{3}\right), \qquad (15)\\
n &= \frac{\omega^2kl}{a^2-l^2}-\alpha\frac{a^2+l^2}{\omega}m+(a^2-l^2)l\left(\frac{\alpha^2}{\omega^2}(\omega^2k+Q^2+G^2)+\frac{\Lambda}{3}\right), \qquad (16)\\
\left(\frac{\omega^2}{a^2-l^2}+3\alpha^2l^2\right)k &= 1+2\alpha\frac{l}{\omega}m-3\alpha^2\frac{l^2}{\omega^2}(Q^2+G^2)-l^2\Lambda. \qquad (17)
\end{aligned}
$$

Now M, Q, G, a, l, α, Λ represent the mass, electric and magnetic charge, angular momentum, NUT parameter, acceleration and cosmological constant, respectively. It is possible to put the metric (5) in the form

$$\mathrm{d}s^2 = \frac{1}{\Omega^2}\left\{\frac{\hat{\mathcal{Q}} - a^2\hat{\mathcal{P}}}{\rho^2}\mathrm{d}t^2 - \frac{2}{\rho^2\Xi}\Big(\hat{\mathcal{Q}}\Theta - a\hat{\mathcal{P}}R\Big)\mathrm{d}t\mathrm{d}\varphi + \frac{1}{\rho^2\Xi^2}\Big(\hat{\mathcal{Q}}\Theta^2 - \hat{\mathcal{P}}R^2)^2\Big)\mathrm{d}\varphi^2 - \frac{\rho^2}{\hat{\mathcal{Q}}}\mathrm{d}r^2 - \frac{\rho^2\sin^2\theta}{\hat{\mathcal{P}}}\mathrm{d}\theta^2\right\}, \tag{18}$$

where

$$\Theta = a\sin^2\theta + 4l\sin^2\theta/2 \qquad\qquad R = r^2 + (a+l)^2. \tag{19}$$

The outer horizon ($r = r_H$) is determined by $\hat{\mathcal{Q}}(r_H) = 0$. The term $1/\Omega^2$ is a conformal factor, so, this term does not contribute to the Hawking radiation, and we can omit it in our future calculations. We will consider matter fields in the Plebański-Demiánski black hole background. As we have both electric and magnetic charge, we can take in consideration that the covariant derivative is [16, 17]

$$D_\mu = \partial_\mu - \mathrm{i}eA_\mu - \mathrm{i}gB_\mu, \tag{20}$$

where e and g are the electric and magnetic charges, respectively. The action is

$$\begin{aligned} S &= \frac{1}{2}\int \mathrm{d}^4x\sqrt{-g}g^{\mu\nu}D_\mu\Phi D_\nu\Phi &(21)\\ &= -\frac{1}{2}\int \mathrm{d}^4x\sqrt{-g}\Phi^*\Big\{g^{tt}\Big[\partial_t^2 - 2\mathrm{i}(eA_t + gB_t)\partial_t - (eA_t + gB_t)^2\Big] + 2g^{t\varphi}\Big[\partial_t\partial_\varphi - \mathrm{i}(eA_\varphi + gB_\varphi)\partial_t \\ &\quad -\mathrm{i}(eA_t + gB_t)\partial_\varphi - (eA_t + gB_t)(eA_\varphi + gB_\varphi)\Big] + g^{\varphi\varphi}\Big[\partial_\varphi^2 - 2\mathrm{i}(eA_\varphi + gB_\varphi)\partial_\varphi - (eA_\varphi + gB_\varphi)^2\Big] \\ &\quad + \frac{1}{\sqrt{-g}}\partial_r(\sqrt{-g}g^{rr}\partial_r) + \frac{1}{\sqrt{-g}}\partial_\theta(\sqrt{-g}g^{\theta\theta}\partial_\theta)\Big\}\Phi, &(22)\end{aligned}$$

where

$$\begin{aligned} g^{tt} &= -\frac{\rho^2(\hat{\mathcal{Q}}\Theta^2 - R^2\hat{\mathcal{P}})}{\hat{\mathcal{P}}\hat{\mathcal{Q}}(R - a\Theta)^2}, & g^{t\varphi} &= -\frac{\rho^2\Xi(\hat{\mathcal{Q}}\Theta - aR\hat{\mathcal{P}})}{\hat{\mathcal{P}}\hat{\mathcal{Q}}(R - a\Theta)^2}, & g^{\varphi\varphi} &= -\frac{\rho^2\Xi^2(\hat{\mathcal{Q}} - a^2\hat{\mathcal{P}})}{\hat{\mathcal{P}}\hat{\mathcal{Q}}(R - a\Theta)^2}, \\ g^{rr} &= -\frac{\hat{\mathcal{Q}}}{\rho^2}, & g^{\theta\theta} &= -\frac{\hat{\mathcal{P}}}{\rho^2\sin^2\theta}, & \sqrt{-g} &= \frac{(R - a\Theta)\,\mathrm{sen}\,\theta}{\Xi}. \end{aligned} \tag{23}$$

Performing the partial wave decomposition of the scalar field Φ in term of the spherical harmonics $\Phi = \sum\limits_{l,m}\phi_{lm}(r,t)Y_{lm}(\theta,\varphi)$ this action can be reduced to a two dimensional effective theory. To do this it is necessary transform to the r_* tortoise coordinate defined by

$$\frac{\mathrm{d}r_*}{\mathrm{d}r} = \frac{R}{\hat{\mathcal{Q}}} = f(r)^{-1}, \tag{24}$$

and considering the region near the horizon, the action (22) can be simplified to

$$S = -\frac{1}{2}\int \mathrm{d}t\mathrm{d}r\frac{R}{\Xi}\phi^*_{l,m}\left\{\frac{R}{\hat{\mathcal{Q}}}\left(\partial_t + \mathrm{i}m\frac{a\Xi}{R} + \mathrm{i}eA_S + \mathrm{i}g\breve{A}_S\right)^2 - \partial_r\frac{\hat{\mathcal{Q}}}{R}\partial_r\right\}\phi_{l,m}; \tag{25}$$

that is, each partial wave of the scalar field can be considered as a $(1+1)$–dimensional complex scalar field in the backgrounds of the dilaton Ψ, metric $\tilde{g}_{\mu\nu}$ and gauge fields $\tilde{A}_\mu$ given by

$$\Psi = \frac{R}{\Xi} = \frac{r^2 + (a+l)^2}{1 + \frac{\Lambda}{3}(a^2 - l^2)}, \tag{26}$$

$$
\begin{aligned}
\tilde{g}_{tt} &= -\frac{\hat{\mathcal{Q}}}{R} = -\frac{(\omega^2 k + Q^2 + G^2) - 2Mr + \epsilon r^2 - 2\alpha \dfrac{n}{\omega} r^3 - \left(\alpha^2 k + \dfrac{\Lambda}{3}\right) r^4}{r^2 + (a+l)^2}, &(27)\\
\tilde{g}_{rr} &= \frac{R}{\hat{\mathcal{Q}}}, &(28)\\
\tilde{A}_{t1} &= -\frac{a\Xi}{R} = -\frac{a\left(1 + \dfrac{\Lambda}{3}(a^2 - l^2)\right)}{r^2 + (a+l)^2}, &(29)\\
\tilde{A}_{t2} &= -A_S = -\frac{Qr - G\dfrac{\Lambda}{3}(l+a)(a^2 - l^2)}{r^2 + (a+l)^2}, &(30)\\
\tilde{A}_{t3} &= -\breve{A}_S = -\frac{Gr + Q\dfrac{\Lambda}{3}(l+a)(a^2 - l^2)}{r^2 + (a+l)^2}, &(31)\\
\tilde{A}_r &= 0; &(32)
\end{aligned}
$$

where m, e, g are the charges of the gage fields $\tilde{A}_{t1}$, $\tilde{A}_{t2}$, $\tilde{A}_{t3}$; respectively.

In this $(1+1)$–dimensional reduction, the effective field theory is based on the observable physics and defined outside the horizon of the black hole. This mean that the ingoing modes are omitted at the horizon doing chiral the theory there. With this, each partial wave become anomalous with respect to gauge and general coordinate symmetries. So, to have gauge invariance and diffeomorphism covariance it is necessary the fluxes of the $U(1)$ gauge current and the energy-momentum tensor cancel the gauge and gravitational anomaly at the horizon, respectively. We will show this procedure in the next section.

3 Anomalies and Hawking radiation

An anomaly in QFT is a conflict between a symmetry from classical action and the quantization. There exist anomalies in global symmetries and gauge symmetries. The gauge current must satisfy the conservation equation $\nabla_\mu J^\mu = 0$. However, near the horizon, the $U(1)$ gauge current satisfies an anomalous equation

$$
\nabla_\mu J^\mu = \frac{\alpha}{4\pi\sqrt{-g}} \epsilon^{\mu\nu} \partial_\mu A_\nu, \tag{33}
$$

where α is the gauge charge of the $U(1)$ gauge field A_ν. So, in the region $r \geq r_H + \epsilon$, as there is no anomaly, the $U(1)$ gauge current must satisfies the conservation equation $\partial_r J^r_{(out)} = 0$ but, near the horizon must satisfies $\partial_r J^r_{(H)} = \dfrac{\alpha \partial_r \tilde{A}_t}{4\pi}$. Thus we can get

$$
\begin{aligned}
J^r_{(out)} &= c_o, & r \geq r_H + \epsilon, \qquad &(34)\\
J^r_{(H)} &= c_H + \frac{1}{4\pi}\Big(\tilde{A}_t(r) - \tilde{A}_t(r_H)\Big), & r_H \leq r \leq r_H + \epsilon; \qquad &(35)
\end{aligned}
$$

where c_o and c_H are integration constants. c_o is the value of current at $r = \infty$ and c_H is the value of consistent current of the outgoing modes at the horizon. To get the values of c_o and c_H, we can use the consistent current

$$
J^\mu = J^\mu_{(out)} \Theta(r - r_H - \epsilon) + J^\mu_{(H)} \left[1 - \Theta(r - r_H - \epsilon)\right], \tag{36}
$$

where

$$
\Theta(r - r_H - \epsilon) = \begin{cases} 1, & r \geq r_H + \epsilon, \\[2ex] 0, & r_H \leq r \leq r_H + \epsilon. \end{cases} \tag{37}
$$

Since we have omitted the ingoing modes near the horizon, this current is only a part of the total current. Under gauge transformation, the variation of the effective action is

$$-\delta W = \int \mathrm{d}t\mathrm{d}r\sqrt{-\tilde{g}}\lambda\nabla_\mu J^\mu, \tag{38}$$

where λ is a gauge parameter. By integration by parts we have

$$-\delta W = \int \mathrm{d}t\mathrm{d}r\lambda\left[\delta(r-r_H-\epsilon)\Big(J^r_{(out)} - J^r_{(H)} + \frac{\alpha}{4\pi}\tilde{A}_t\Big) + \partial_t\Big(\frac{\alpha}{4\pi}\tilde{A}_t\left[1-\Theta(r-r_H-\epsilon)\right]\Big)\right]. \tag{39}$$

The total effective action must be gauge invariant. So, the last term would be vanish by quantum effects of the classically irrelevant ingoing modes. The coefficient of the delta function would be canceled also, and we get

$$c_o = c_H - \frac{\alpha}{4\pi}\tilde{A}_t(r_H). \tag{40}$$

To ensure the regularity requirement at the horizon, the covariant current must also vanish there. Since the covariant current is $\tilde{J}^r = J^r + \frac{\alpha}{4\pi}\tilde{A}_t\left[1-\Theta(r-r_H-\epsilon)\right]$, the condition at the horizon $\tilde{J}^r = 0$ determines the flux of the $U(1)$ gauge current as

$$c_o = -\frac{\alpha}{2\pi}\tilde{A}_t(r_H). \tag{41}$$

We have three gauge charges for a Plebański-Demiański black hole; thus, the $U(1)$ gauge charge flux, the electric current flux and the magnetic current flux are determined by

$$f_m = -\frac{m}{2\pi}\tilde{A}_t(r_H) = \frac{m^2a\Xi + me\Big(Qr_H - G(a+l)(\Xi-1)\Big) + mg\Big(Gr_H + Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_H^2+(a+l)^2\Big)}, \tag{42}$$

$$f_e = -\frac{e}{2\pi}\tilde{A}_t(r_H) = \frac{mea\Xi + e^2\Big(Qr_H - G(a+l)(\Xi-1)\Big) + eg\Big(Gr_H + Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_H^2+(a+l)^2\Big)}, \tag{43}$$

$$f_g = -\frac{g}{2\pi}\tilde{A}_t(r_H) = \frac{mga\Xi + ge\Big(Qr_H - G(a+l)(\Xi-1)\Big) + g^2\Big(Gr_H + Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_H^2+(a+l)^2\Big)}. \tag{44}$$

The anomalies in global symmetries are theoretical inconsistences for this reason their cancellation give important restrictions [18, 19]. A gravitational anomaly is a gauge anomaly in general covariance, making non-conservative the energy-momentum tensor. This anomaly can only happen in theories with chiral matter acopled to gravity in a space-time $4k+2$–dimensional, being k an integer. This chiral matter can be a fermion and also can be a $2k$–form with an (anti–) autodual field. An important situation is the $(1+1)$–dimensional scalar autodual field; this field obeys

$$\partial_a\phi = \epsilon_{ab}\partial^b\phi, \tag{45}$$

that is, it has only modes are moving to right, so it is chiral. The anomaly is

$$\nabla_a T^a_b = \frac{1}{\sqrt{-g}}\partial_a N^a_b = \frac{1}{96\pi\sqrt{-g}}\epsilon^{cd}\partial_d\partial_a\Gamma^a_{bc}; \tag{46}$$

that is, the energy-momentum tensor is not conserved in a curved Space-Time. As discussed previously, the effective field theory is defined outside the event horizon. In the region $r \geq r_H + \epsilon$ there is an effective background gauge potential, but without anomaly, so the energy-momentum tensor satisfies the modified conservation equation $\partial_r T^r_{t(out)} = c_o\partial_r\tilde{A}_t$. Near the horizon $r_H \leq r \leq r_H + \epsilon$ the energy-momentum tensor exhibits an anomaly and satisfies the Ward identity, that is $\partial_r T^r_{t(H)} = J^r_{(H)}\partial_r\tilde{A}_t + \tilde{A}_t\partial_r J^r_{(H)} + \partial_r N^r_t$, where

$$N^r_t = \frac{1}{192\pi}\frac{\hat{\mathcal{Q}}\mathcal{M} + \mathcal{C}^2}{R^4}, \tag{47}$$

and

$$\begin{aligned}\mathcal{C} &= \hat{\mathcal{Q}}'R - 2r\hat{\mathcal{Q}}, &(48)\\ &= 2M\Big(r^2-(a+l)^2\Big)+2\Big(\epsilon(a+l)^2-\omega^2k-Q^2-G^2\Big)r-2\alpha\frac{n}{\omega}\Big(r^2+3(a+l)^2\Big)r^2\\ &\quad -2\Big(\alpha^2k+\frac{\Lambda}{3}\Big)\Big(r^2+2(a+l)^2\Big)r^3, &(49)\\ \mathcal{M} &= \mathcal{C}'R-2R'\mathcal{C}, &(50)\\ &= 8M(a+l)^2r-2\Big(\epsilon(a+l)^2-\omega^2k-Q^2-G^2\Big)\Big(3r^2-(a+l)^2\Big)+2\alpha\frac{n}{\omega}\Big(5(a+l)^2r^3-3(a+l)^4r\Big)\\ &\quad -2\Big(\alpha^2k+\frac{\Lambda}{3}\Big)\Big(r^6+3(a+l)^2r^4+6(a+l)^4r^2\Big). &(51)\end{aligned}$$

We can use the energy-momentum tensor

$$T^r_t = T^r_{t(out)}\Theta(r-r_H-\epsilon)+T^r_{t(H)}\left[1-\Theta(r-r_H-\epsilon)\right], \tag{52}$$

which combines contributions from these two regions.

If a classical action $S[\Phi, g_{\mu\nu}]$ is quantized, we get

$$\mathcal{W}[g_{\mu\nu}] = -\mathrm{i}\ln\left(\int\mathcal{D}\Phi e^{\mathrm{i}S[\Phi,g_{\mu\nu}]}\right); \tag{53}$$

and, under general coordinate transformation $x^\mu \longrightarrow x^\mu - \lambda^\mu$, the variation of the effective action become

$$\begin{aligned}-\delta_\lambda\mathcal{W} &= \int \mathrm{d}t\mathrm{d}r\lambda^\nu\nabla_\mu T^\mu_\nu &(54)\\ &= \int \mathrm{d}t\mathrm{d}r\lambda^t\Big[c_o\partial_r\tilde{A}_t+\partial_r\Big(\frac{\alpha}{4\pi}\tilde{A}^2_t+N^r_t\Big)\left[1-\Theta(r-r_H-\epsilon)\right]\\ &\qquad +\Big(T^r_{t(out)}-T^r_{t(H)}+\frac{\alpha}{4\pi}\tilde{A}^2_t+N^r_t\Big)\delta(r-r_H-\epsilon)\Big]; &(55)\end{aligned}$$

where the relation $J^r_{(H)} = c_o + \frac{\alpha}{4\pi}\tilde{A}_t$ has been used. The first term of the effective action is the classical effect of the background electric field. The second term is cancelled by the quantum effect of the classically irrelevant ingoing modes. The third one would be cancelled if we demand the covariance of effective action under the diffeomorphism transformation, this let us

$$a_o = a_H + \frac{\alpha}{4\pi}\tilde{A}^2_t(r_H) - N^r_t(r_H), \tag{56}$$

where

$$\begin{aligned}a_o &= T^r_{t(out)} - c_o\tilde{A}_t, &(57)\\ a_H &= T^r_{t(H)} - \int_{r_H}^{r}\mathrm{d}r\partial_r\left[c_o\tilde{A}_t+\frac{\alpha}{4\pi}\tilde{A}^2_t+N^r_t\right], &(58)\end{aligned}$$

are the values of the energy flow at infinity and at the horizon, respectively. To ensure the regularity requirement at the horizon, the covariant energy-momentum tensor must also vanish there. Since the energy-momentum tensor is $\tilde{T}^r_t = T^r_t + \frac{1}{192\pi}\frac{\hat{\mathcal{Q}}\mathcal{M}-2\mathcal{C}^2}{R^4}$, the condition at the horizon $\tilde{T}^r_t = 0$ carry us

$$a_H = 2N^r_t(r_H) = \frac{\kappa^2}{24\pi}; \tag{59}$$

where

$$
\begin{aligned}
\kappa &= \frac{1}{2}\frac{\mathcal{C}}{R^2}\Big|_{r=r_H}, \qquad (60)\\
&= \frac{M\Big(r_H^2-(a+l)^2\Big)+r_H\Big(\epsilon(a+l)^2-\omega^2 k-Q^2-G^2\Big)-\alpha\frac{n}{\omega}\Big(r_H^2+3(a+l)^2\Big)r_H^2}{r_H^2+(a+l)^2}\\
&\quad -\frac{\Big(\alpha^2 k+\frac{\Lambda}{3}\Big)\Big(r_H^2+2(a+l)^2\Big)r_H^3}{r_H^2+(a+l)^2}, \qquad (61)
\end{aligned}
$$

is the surface gravity of the black hole. Then, the flux of the energy-momentum tensor required to restore general coordinate covariance at quantum level in the effective field theory is

$$
\begin{aligned}
a_o &= N_t^r(r_H)+\frac{\alpha}{4\pi}\tilde{A}_t^2(r_H), \qquad (62)\\
&= \frac{1}{4\pi}\left(\frac{ma\Xi+e\Big(Qr_H-G(a+l)(\Xi-1)\Big)+g\Big(Gr_H+Q(a+l)(\Xi-1)\Big)}{r_H^2+(a+l)^2}\right)^2+\frac{\pi}{12}T_h^2 \qquad (63)
\end{aligned}
$$

where $T_h=\dfrac{\kappa}{2\pi}$ is Hawking temperature of the black hole. The Hawking radiant spectrum of the Plebański-Demiański black hole is given by the Planck distribution

$$
N_{\pm m,\pm e,\pm g}=\frac{1}{exp\Big(\frac{\omega\mp m\tilde{A}_{t_1}\mp e\tilde{A}_{t_2}\mp g\tilde{A}_{t_3}}{T_h}\Big)+1}. \tag{64}
$$

From the distribution, the angular momentum flux (that is, the $U(1)$ gauge current flux), the electric current flux and the magnetic current flux can be obtained as

$$
\begin{aligned}
F_m &= m\int_0^\infty \frac{1}{2\pi}\Big[N_{m,e,g}(\omega)-N_{-m,-e,-g}(\omega)\Big]\mathrm{d}\omega \qquad (65)\\
&= \frac{m^2a\Xi+me\Big(Qr_H-G(a+l)(\Xi-1)\Big)+mg\Big(Gr_H+Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_H^2+(a+l)^2\Big)}, \qquad (66)
\end{aligned}
$$

$$
\begin{aligned}
F_e &= e\int_0^\infty \frac{1}{2\pi}\Big[N_{m,e,g}(\omega)-N_{-m,-e,-g}(\omega)\Big]\mathrm{d}\omega \qquad (67)\\
&= \frac{mea\Xi+e^2\Big(Qr_H-G(a+l)(\Xi-1)\Big)+eg\Big(Gr_H+Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_H^2+(a+l)^2\Big)}, \qquad (68)
\end{aligned}
$$

$$
\begin{aligned}
F_e &= e\int_0^\infty \frac{1}{2\pi}\Big[N_{m,e,g}(\omega)-N_{-m,-e,-g}(\omega)\Big]\mathrm{d}\omega \qquad (69)\\
&= \frac{mga\Xi+ge\Big(Qr_H-G(a+l)(\Xi-1)\Big)+g^2\Big(Gr_H+Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_H^2+(a+l)^2\Big)}; \qquad (70)
\end{aligned}
$$

and the energy-momentum tensor current is

$$
F_H = \int_0^\infty \frac{\omega}{2\pi}\Big[N_{m,e,g}(\omega)-N_{-m,-e,-g}(\omega)\Big]\mathrm{d}\omega \tag{71}
$$

$$= \frac{1}{4\pi}\left(\frac{ma\Xi + e\Big(Qr_H - G(a+l)(\Xi-1)\Big) + g\Big(Gr_H + Q(a+l)(\Xi-1)\Big)}{r_H^2 + (a+l)^2}\right)^2 + \frac{\pi}{12}T_h^2. \quad (72)$$

Comparing equations (42–44,63) with equations (66,68,70,72), we can conclude that the fluxes of the $U(1)$ gauge current, electric current, magnetic current and the energy-momentum tensor required to cancel gauge or gravitational anomalies at horizon are identical to that of Hawking radiation.

4 Hawking radiation at Cosmological horizon

Near the Cosmological horizon (CH), we must take in consideration ingoing modes. Inside the CH ($r_c - \epsilon < r < r_c$), the modes need to satisfy the equation $\partial_r J^r_{(c)} = -\frac{\alpha\partial_r\tilde{A}_t}{4\pi}$, but in $r < r_c - \epsilon$ there is no anomaly, so we have a conservation equation $\partial_r J^r_{(c-out)} = 0$. We can get from this equations, the consistent current

$$J^r = J^r_{(c-out)}\Theta(r_c - \epsilon - r) + J^r_{(c)}[1 - \Theta(r_c - \epsilon - r)]. \quad (73)$$

Under gauge transformation, the action variation become

$$-\delta W = \int \mathrm{d}t\mathrm{d}r\lambda\left[\delta(r_c - \epsilon - r)\Big(J^r_{(c)} - J^r_{(c-out)} + \frac{\alpha}{4\pi}\tilde{A}_t\Big) + \partial_t\Big(\frac{\alpha}{4\pi}\tilde{A}_t\left[1 - \Theta(r_c - \epsilon - r)\right]\Big)\right]. \quad (74)$$

As the covariant current must vanish at the CH, we obtain the $U(1)$ gauge charge flux, the electric current flux and the magnetic current flux at the CH, are determined by

$$\tilde{f}_m = \frac{m}{2\pi}\tilde{A}_t(r_c) = -\frac{m^2a\Xi + me\Big(Qr_c - G(a+l)(\Xi-1)\Big) + mg\Big(Gr_c + Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_c^2 + (a+l)^2\Big)}, \quad (75)$$

$$\tilde{f}_e = \frac{e}{2\pi}\tilde{A}_t(r_c) = -\frac{mea\Xi + e^2\Big(Qr_c - G(a+l)(\Xi-1)\Big) + eg\Big(Gr_c + Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_c^2 + (a+l)^2\Big)}, \quad (76)$$

$$\tilde{f}_g = \frac{g}{2\pi}\tilde{A}_t(r_c) = -\frac{mga\Xi + ge\Big(Qr_c - G(a+l)(\Xi-1)\Big) + g^2\Big(Gr_c + Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_c^2 + (a+l)^2\Big)}. \quad (77)$$

At similar form, in $r_c - \epsilon < r < r_c$, the energy-momentum tensor satisfies the Ward identity $\partial_r T^r_{t(c)} = J^r_{(c)}\partial_r\tilde{A}_t + \tilde{A}_t\partial_r J^r_{(c)} - \partial_r N^r_t$, and, in $r < r_c - \epsilon$, the energy-momentum satisfies $\partial_r T^r_{t(c-out)} = \tilde{a}_o\partial_r\tilde{A}_t$ If we use the energy-momentum tensor

$$T^r_t = T^r_{t(c-out)}\Theta(r_c - \epsilon - r) + T^r_{t(c)}[1 - \Theta(r_c - \epsilon - r)], \quad (78)$$

after general coordinate transformation, the variation of the effective action become

$$\begin{aligned} -\delta_\lambda\mathcal{W} &= \int \mathrm{d}t\mathrm{d}r\lambda^t\Big[\tilde{c}_o\partial_r\tilde{A}_t + \partial_r\Big(\frac{\alpha}{4\pi}\tilde{A}_t^2 + N^r_t\Big)\left[1 - \Theta(r_c - \epsilon - r)\right] \\ &\quad + \Big(T^r_{t(c)} - T^r_{t(c-out)} + \frac{\alpha}{4\pi}\tilde{A}_t^2 + N^r_t\Big)\delta(r_c - \epsilon - r)\Big]; \end{aligned} \quad (79)$$

and, as the covariant energy-momentum tensor must vanish at the CH, the flux of the energy-momentum tensor required to restore general coordinate covariance at quantum level in the effective field theory is

$$\tilde{a}_o = -\frac{1}{4\pi}\left(\frac{ma\Xi + e\Big(Qr_c - G(a+l)(\Xi-1)\Big) + g\Big(Gr_c + Q(a+l)(\Xi-1)\Big)}{r_c^2 + (a+l)^2}\right)^2 - \frac{\pi}{12}T_c^2; \quad (80)$$

where $T_c = -\frac{1}{4}\frac{\mathcal{C}}{R^2}\Big|_{r=r_c}$ is the temperature of black hole.

The Hawking radiant spectrum of the Plebański-Demiański black hole at CH is given by the Planck distribution

$$N_{\pm m,\pm e,\pm g} = -\frac{1}{exp\Big(\dfrac{\omega \mp m\tilde{A}_{t_1} \mp e\tilde{A}_{t_2} \mp g\tilde{A}_{t_3}}{T_h}\Big) + 1}. \tag{81}$$

From the distribution, the $U(1)$ gauge current flux, the electric current flux and the magnetic current flux can be obtained as

$$\begin{aligned} F_m &= m\int_0^\infty \frac{1}{2\pi}\Big[N_{m,e,g}(\omega) - N_{-m,-e,-g}(\omega)\Big]\mathrm{d}\omega & (82)\\ &= -\frac{m^2a\Xi + me\Big(Qr_H - G(a+l)(\Xi-1)\Big) + mg\Big(Gr_H + Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_H^2 + (a+l)^2\Big)}, & (83)\\ F_e &= e\int_0^\infty \frac{1}{2\pi}\Big[N_{m,e,g}(\omega) - N_{-m,-e,-g}(\omega)\Big]\mathrm{d}\omega & (84)\\ &= -\frac{mea\Xi + e^2\Big(Qr_H - G(a+l)(\Xi-1)\Big) + eg\Big(Gr_H + Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_H^2 + (a+l)^2\Big)}, & (85)\\ F_g &= g\int_0^\infty \frac{1}{2\pi}\Big[N_{m,e,g}(\omega) - N_{-m,-e,-g}(\omega)\Big]\mathrm{d}\omega & (86)\\ &= -\frac{mga\Xi + ge\Big(Qr_H - G(a+l)(\Xi-1)\Big) + g^2\Big(Gr_H + Q(a+l)(\Xi-1)\Big)}{2\pi\Big(r_H^2 + (a+l)^2\Big)}, & (87) \end{aligned}$$

and the energy-momentum tensor current is

$$\begin{aligned} F_H &= \int_0^\infty \frac{\omega}{2\pi}\Big[N_{m,e,g}(\omega) - N_{-m,-e,-g}(\omega)\Big]\mathrm{d}\omega & (88)\\ &= -\frac{1}{4\pi}\left(\frac{ma\Xi + e\Big(Qr_H - G(a+l)(\Xi-1)\Big) + g\Big(Gr_H + Q(a+l)(\Xi-1)\Big)}{r_H^2 + (a+l)^2}\right)^2 - \frac{\pi}{12}T_h^2. & (89) \end{aligned}$$

Comparing equations (75–77,80) with equations (83,85,87,89), we can conclude that the fluxes of the $U(1)$ gauge current, electric current, magnetic current and the energy-momentum tensor required to cancel gauge or gravitational anomalies at cosmological horizon are identical to that of Hawking radiation.

5 Limit cases

Black holes without NUT parameter

An interesting case is obtained if there is not NUT parameter ($l = 0$), the PD metric (5) is reduced to

$$\begin{aligned}
&\frac{\omega^2}{a^2}k = 1, \qquad \Longrightarrow \qquad \begin{cases} k = 1, \\ \omega = a, \\ \epsilon = 1 - \alpha^2(a^2 + Q^2 + G^2) - \dfrac{\Lambda}{3}a^2, \\ n = -\alpha a M, \end{cases} \\
&a_3 = 2\alpha m, \qquad a_4 = -\alpha^2(a^2 + Q^2 + G^2) - \frac{\Lambda}{3}a^2, \\
&\Omega = 1 - \alpha r\cos\theta, \qquad \rho^2 = r^2 + a^2\cos^2\theta, \\
&\mathcal{P} = \sin^2\theta\left[1 - 2\alpha M\cos\theta + \left(\alpha^2(a^2 + Q^2 + G^2) + \frac{\Lambda}{3}a^2\right)\cos^2\theta\right], \\
&\hat{\mathcal{Q}} = (r^2 + a^2 + Q^2 + G^2 - 2Mr)(1 - \alpha^2 r^2) - \frac{\Lambda}{3}r^2(a^2 + r^2);
\end{aligned} \tag{90}$$

and at this form the accelerated Kerr-Newman-de Sitter with magnetic monopole black hole metric is obtained. If we considerate $\alpha = 0$ and $G = 0$, we will get

$$\hat{\mathcal{Q}} = r^2 + a^2 + Q^2 - 2Mr - \frac{\Lambda}{3}r^2(r^2 + a^2), \tag{91}$$

and the metric become the Kerr-Newman-de Sitter one

$$\begin{aligned}
\mathrm{d}s^2 \;=\;& \frac{r^2 - 2Mr + a^2 + Q^2 + -\frac{\Lambda}{3}r^2(r^2 + a^2)}{r^2 + a^2\cos^2\theta}\left(\mathrm{d}t - \frac{a\sin^2\theta}{1 + \frac{\Lambda}{3}a^2}\mathrm{d}\varphi\right)^2 - \frac{r^2 + a^2\cos^2\theta}{1 + \frac{\Lambda}{3}a^2\cos^2\theta}\mathrm{d}\theta^2 \\
& - \frac{r^2 + a^2\cos^2\theta}{r^2 - 2Mr + a^2 + Q^2 - \frac{\Lambda}{3}r^2(r^2 + a^2)}\mathrm{d}r^2 - \frac{\sin^2\theta\left(1 + \frac{\Lambda}{3}a^2\cos^2\theta\right)}{r^2 + a^2\cos^2\theta}\left(a\mathrm{d}t - \frac{r^2 + a^2}{1 + \frac{\Lambda}{3}a^2}\mathrm{d}\varphi\right)^2 .
\end{aligned} \tag{92}$$

As there not exists magnetic charge, the value of $g = 0$, and, taking the value of $\hat{\mathcal{Q}}$ (ec. 91), we get the gauge field

$$\tilde{A}_t = -\frac{ma\left(1 + \frac{\Lambda}{3}a^2\right) + eQr}{r^2 + a^2}; \tag{93}$$

and the $U(1)$ gauge charge flux, electric current flux, energy-momentum flux and Hawking temperature are

$$f_m \;=\; -\frac{m}{2\pi}\tilde{A}_t(r_H) = \frac{m^2 a\left(1 + \frac{\Lambda}{3}a^2\right) + meQr_H}{2\pi\left(r_H^2 + a^2\right)}, \tag{94}$$

$$f_e \;=\; -\frac{e}{2\pi}\tilde{A}_t(r_H) = \frac{mea\left(1 + \frac{\Lambda}{3}a^2\right) + e^2Qr_H}{2\pi\left(r_H^2 + a^2\right)}, \tag{95}$$

$$a_o \;=\; \frac{1}{4\pi}\left(\frac{ma\left(1 + \frac{\Lambda}{3}a^2\right) + eQr_H}{r_H^2 + a^2}\right)^2 + \frac{\pi}{12}T_h^2, \tag{96}$$

$$T_h \;=\; \frac{M(r_H^2 - a^2) - r_H Q^2 - \frac{\Lambda}{3}r_H(r_H^2 + a^2)^2}{2\pi(r_H^2 + a^2)}, \tag{97}$$

respectively; this values corresponding to the values obtained in [11].

Non accelerated black holes

Another interesting case is obtained if the black hole is not accelerated ($\alpha = 0$). The PD metric (5) is reduced to

$$\begin{aligned}
&\frac{\omega^2 k}{a^2 - l^2} = 1 - l^2\Lambda, \qquad \Longrightarrow \qquad \begin{cases} \epsilon = 1 - \dfrac{\Lambda}{3}(a^2 + 6l^2), \\ n = l + \dfrac{\Lambda}{3} l (a^2 - 4l^2), \end{cases} \\
&a_3 = -4\frac{\Lambda}{3} a l, \qquad a_4 = -\frac{\Lambda}{3} a^2, \\
&\Omega = 1 \qquad \rho^2 = r^2 + (l + a\cos\theta)^2, \\
&\mathcal{P} = \sin^2\theta \left(1 + 4\frac{\Lambda}{3} a l \cos\theta + \frac{\Lambda}{3} a^2 \cos^2\theta\right), \\
&\hat{\mathcal{Q}} = r^2 + a^2 + Q^2 + G^2 - l^2 - 2Mr - \frac{\Lambda}{3}\Big(r^4 + r^2(a^2 + 6l^2) + 3l^2(a^2 - l^2)\Big);
\end{aligned} \tag{98}$$

and we obtain the NUT-Kerr-Newman-de Sitter with magnetic monopole black hole metric. If we have no magnetic monopole, we get

$$\hat{\mathcal{Q}} = r^2 + a^2 + Q^2 + -l^2 - 2Mr - \frac{\Lambda}{3}\Big(r^4 + r^2(a^2 + 6l^2) + 3l^2(a^2 - l^2)\Big), \tag{99}$$

and the metric become the NUT-Kerr-Newman-de Sitter one

$$\begin{aligned}
\mathrm{d}s^2 \quad = \quad & \frac{r^2 + a^2 + Q^2 - l^2 - 2Mr - \frac{\Lambda}{3}\Big(r^4 + r^2(a^2 + 6l^2) + 3l^2(a^2 - l^2)\Big)}{r^2 + (l + a\cos\theta)^2} \left(\mathrm{d}t - \frac{a\sin^2\theta + 4l\sin^2\frac{\theta}{2}}{1 + \frac{\Lambda}{3}(a^2 - l^2)}\mathrm{d}\varphi\right)^2 \\
& - \frac{\sin^2\theta\left(1 + 4\frac{\Lambda}{3} a l \cos\theta + \frac{\Lambda}{3} a^2\cos^2\theta\right)}{r^2 + (l + a\cos\theta)^2}\left(a\mathrm{d}t - \frac{r^2 + (a + l)^2}{1 + \frac{\Lambda}{3}(a^2 - l^2)}\mathrm{d}\varphi\right)^2 - \frac{r^2 + (l + a\cos\theta)^2}{1 + 4\frac{\Lambda}{3} a l\cos\theta + \frac{\Lambda}{3} a^2\cos^2\theta}\mathrm{d}\theta^2 \\
& - \frac{r^2 + (l + a\cos\theta)^2}{r^2 + a^2 + Q^2 - l^2 - 2Mr - \frac{\Lambda}{3}\Big(r^4 + r^2(a^2 + 6l^2) + 3l^2(a^2 - l^2)\Big)}\mathrm{d}r^2.
\end{aligned} \tag{100}$$

As there not exists magnetic charge, the value of $g = 0$, and, taking the value of $\hat{\mathcal{Q}}$ (ec. 99), we get the gauge field

$$\tilde{A}_t = -\frac{ma\Big(1 + \frac{\Lambda}{3}(a^2 - l^2)\Big) + eQr}{r^2 + (a + l)^2}; \tag{101}$$

and the $U(1)$ gauge charge flux, electric current flux, energy-momentum flux and Hawking temperature are

$$f_m \quad = \quad \frac{m^2 a\Big(1 + \frac{\Lambda}{3}(a^2 - l^2)\Big) + meQr_H}{2\pi\Big(r_H^2 + (a + l)^2\Big)}, \tag{102}$$

$$f_e \quad = \quad \frac{mea\Big(1 + \frac{\Lambda}{3}(a^2 - l^2)\Big) + e^2Qr_H}{2\pi\Big(r_H^2 + (a + l)^2\Big)}, \tag{103}$$

$$a_o \quad = \quad \frac{1}{4\pi}\left(\frac{ma\Big(1 + \frac{\Lambda}{3}(a^2 - l^2)\Big) + eQr_H}{r_H^2 + (a + l)^2}\right)^2 + \frac{\pi}{12}T_h^2, \tag{104}$$

$$T_h \quad = \quad \frac{M\Big(r_H^2-(a+l)^2\Big)+r_H\left[2l(a+l)-Q^2-\dfrac{\Lambda}{3}\left(\Big[r_H^2+(a+l)^2\Big]^2+2l(a+l)(4l^2-a^2)\right)\right]}{2\pi\Big(r_H^2+(a+l)^2\Big)}, \qquad (105)$$

respectively; this values are founded in [13]. It is important to observe that expresions (99–105) reduced to (91–97) when $l=0$.

6 Conclusions

After the Robinson-Wilczek article [7], where it was showed how to get the Hawking radiation for a Schwarzschild black hole via anomalies, different groups started to show the Hawking radiation for more general black holes. We have showed Hawking radiation for a Plebański-Demiański black hole in this article, that is, for the most general D–type metric. It is important to remark that any particular combination of parameters included in the PD black hole (mass, electric and magnetic charge, angular momentum, NUT parameter, acceleration and cosmological constant) can be obtained from our calculations and we have showed this obtaining special cases previously reported.

With this generalization, we can be sure that near tho horizon of a black hole, there is quantum vacuum fluctuation effect and virtual particle pairs can appear. The antiparticles enter the event horizon of the black hole and become ingoing modes. At this form, the theory become anomalous. The anomaly can be cancelled originating as consequence the Hawking radiation.

References

[1] Ya.B. Zeldovich, A.A. Starobinsky, *Particle creation and vacuum polarization in an anisotropic gravitational field*, Zh.E.T.F. **61** (1971) 2161, [JETP **34** (1972) 1159].

[2] S.W. Hawking, *Black hole explosions?*, Nature **248** (1974) 30.

[3] S.W. Hawking, *Particle creation by black holes*, Commun. Math. Phys. **43** (1975) 199.

[4] G. Gibbons, S.W. Hawking, *Action integrals and partition functions in quantum gravity*, Phys. Rev. D **15** (1977) 2752.

[5] A. Strominger, C. Vafa, *Microscopic origin of the Bekenstein-Hawking entropy*, Phys. Lett. B **379** (1996) 99. [hep-th/9601029].

[6] A. Peet, *TASI lectures on black holes in string theory.* [hep-th/0008241].

[7] S.P. Robinson, F. Wilczek, *Relationship between Hawking radiation anf gravitational anomalies*, Phys. Rev. Lett. **95** (2005) 011303.

[8] S. Iso, H. Umetsu, F. Wilczek, *Hawking radiation from charged black holes via gauge and gravitational anomalies*, Phys. Rev. Lett. **96** (2006) 151302. [hep-th/0602146].

[9] S. Iso, H. Umetsu, F. Wilczek, *Anomalies, Hawking radiation and regularity in rotating black holes*, Phys. Rev. D **74** (2006) 044017. [hep-th/0606018].

[10] K. Murata, J. Soda, *Hawking radiation from rotating black holes and gravitational anomalies*, Phys. Rev. D **74** (2006) 044018. [hep-th/0606069].

[11] Q.Q. Jiang, S.Q. Wu, *Hawking radiation from rotating black holes in anti-de Sitter spaces via gauge and gravitational anomalies*, Phys. Lett. B **647** (2007) 200. [hep-th/0701002].

[12] S. Iso, T. Morita, H. Umetsu, *Quantum anomalies at Horizon and Hawking radiations in Myers-Perry black holes*, JHEP **04** (2007) 068. [hep-th/0612286].

[13] K. Lin, S.W. Chen, S.Z. Yang, *Anomalies and Hawking radiation of NUT-Kerr-Newman de Sitter black hole*, Int. J. Theor. Phys. **47** (2008) 2453.

[14] J.F. Plebański, M. Demiański, *Rotating charged and uniformly accelerating mass in general relativity*, Annals Phys. **98** (1976) 98.

[15] J.B. Griffiths, J. Podolský, *A new look at the Plebański-Demiański family of solutions*, Int. J. Mod. Phys. **15** (2006) 335.

[16] D. Zwanziger, *Local-lagrangian field theory of electric and magnetic charges*, Phys. Rev. D **3** (1971) 880.

[17] D. Singleton, *Magnetic charge as a "Hidden" gauge symmetry*, Int. J. Theor. Phys. **34** (1995) 37. [hep-th/9701044].

[18] R. Bertlmann, *Anomalies in Quantum Field Theory*, Oxford: Oxford Univ. Press, 1996.

[19] A. Bilal, *Lectures on anomalies*. arXiv:0802.0634 [hep-th].